

Award Number: DAMD17-00-1-0441

TITLE: Isolation of Factors that Disrupt Critical
Protein/Protein Interactions within the Telomerase
Holoenzyme for Use in Breast Cancer Therapies

PRINCIPAL INVESTIGATOR: Michael A. White, Ph.D.

CONTRACTING ORGANIZATION: University of Texas Southwestern Medical
Center at Dallas
Dallas, Texas 75390-9105

REPORT DATE: June 2003

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20031104 093

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 074-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE Jun 2003	3. REPORT TYPE AND DATES COVERED Final (1 Jun 00 - 31 May 03)		
4. TITLE AND SUBTITLE Isolation of Factors that Disrupt Critical Protein/Protein Interactions within the Telomerase		5. FUNDING NUMBERS DAMD17-00-1-0441		
6. AUTHOR(S) Michael A. White, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Texas Southwestern Medical Center at Dallas Dallas, Texas 75390-9105 E-Mail: Michael.white@utsouthwestern.edu		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 Words) Telomerase activity is required to maintain telomere integrity on chromosomes of proliferating cells and thus is critically involved in regulating cellular replicative lifespan. Telomerase is repressed in most adult somatic cells, and activation of telomerase activity is an early event associated with tumor progression. Expression of telomerase is sufficient to greatly prolong proliferative lifespan of human cells in culture. Because telomerase activity is not required to maintain viability of post-mitotic somatic cells, but is required to maintain the proliferative capacity of tumor cells, telomerase is an ideal target for anti-cancer therapies. Here we have produced mammalian expression vectors containing PolIII-driven short-hairpin siRNA precursors targeting hTERT mRNA. We show that these vectors dramatically repress telomerase activity when delivered to telomerase positive immortal human tumor cells, resulting in dramatic telomere shortening and a limited replicative life-span in culture.				
14. SUBJECT TERMS			15. NUMBER OF PAGES 5	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

Table of Contents

Cover.....	1
SF 298.....	2
Introduction.....	3
Body.....	3
Key Research Accomplishments.....	4
Reportable Outcomes.....	5
Conclusions.....	5
References.....	
Appendices.....	

Introduction:

The goal of our grant is to apply our knowledge of the functional assembly of telomerase to design novel inhibitors of telomerase activity as a mechanism of limiting the proliferative potential of breast tumor cells *in vivo*. The telomerase holoenzyme is a multi-subunit complex consisting of a reverse transcriptase catalytic core (hTERT) and an RNA template component (hTR) together with other known and unknown accessory factors. It has recently been demonstrated that small interfering RNA duplexes (siRNAs) can effectively and specifically mediate degradation of mRNAs in human cells. We have assessed the efficacy of siRNA mediated gene silencing in normal and tumor-derived human mammary epithelial cells. We find specific silencing of gene products can be achieved at high efficiency in these cells. Therefore, we are developing a system for delivery of siRNAs targeting hTERT and hTR.

Body:

Our original approach to inhibition of telomerase was through isolation of peptide aptamers that disrupt telomerase assembly by disrupting association of hTERT with p23. As described in last year's progress report, we successfully isolated a pool of peptide aptamers that bind hTERT such that they block association with p23 (task 1). Unfortunately, we have had great difficulty introducing these peptides into cells at sufficient concentrations to compete with native p23 (a relatively abundant protein) (task 2). While there are a number of strategies that can be employed to find a solution to this problem, we have elected to alter our approach to take advantage of recent technological developments in the field of RNAi. This modification of our work plan was approved following last year's progress report.

This new approach has been very successful, and as described below, accomplishes the goals of the original proposal.

Our progress with this approach is as follows: 1) Synthetic siRNA duplexes were synthesized targeting unique 21 nucleotide sequences in the hTERT mRNA and hTR. These were introduced into telomerase positive breast cancer derived cell lines to identify siRNAs that effectively reduced telomerase activity. Anti-hTERT siRNAs with the capacity to dramatically and specifically inhibit hTERT expression were identified. No siRNAs designed to target hTR showed any efficacy *in vivo*. 2) Replication defective retroviral vectors were constructed that express short hairpin RNAs from polIII promoters corresponding to the effective synthetic siRNAs targeting hTERT. Viral particles were produced and used to infect breast cancer cell lines. 3) 12 cell lines were cloned following infection. Of these, three lines showed reduced TRAP activity (Fig. 1). Clone C8 failed to expand further- presumably due to replicative senescence. Clones C2 and C5 were expanded for a total of 15 population doublings, then telomere length was assessed using standard TRF assay (Fig. 1). Both cell lines displayed dramatically shortened telomeres relative to the parentals and parallel clonal control lines. The telomerase impaired clones failed to proliferate beyond approximately 10 additional population doublings.




Figure 1. anti-hTERT shRNA inhibits telomerase activity and telomere elongation in human tumor cells. Two stable clones expressing anti-hTERT shRNA (C2, C5), a control clone (expressing anti-RalB siRNA) and a pooled population of parental cells are shown. Left panel: TRAP activity from equal cell equivalents. Right panel: telomere restriction fragment analysis.

Current focus: 1) We are constructing adenoviral vectors based on the successful retroviral vectors described in 2 above. The reason for this is that we have discovered that polIII-mediated expression of shRNAs is far more effective when using episomal vectors than when integrated into the genome. 2) We are evaluating the efficacy of systemic delivery of synthetic siRNA-related compounds targeting telomerase in mouse xenograft model systems.

Key Research accomplishments:

- isolation of a panel of peptide aptamers that competitively association with the p23 binding domain of hTERT
- validation of siRNA technology in human mammary epithelial cells.
- generation of retroviral expression vectors for stable expression of anti-hTERT shRNA.
- validation of anti-hTERT shRNA as a mechanism to limit replicative lifespan of human tumor cells.

Reportable outcomes: These observations are currently being prepared for publication. The anti-hTERT shRNA retroviral vectors will be made freely available to the research community.

Conclusions:

The reagents that are being developed will be useful in examining the biological consequences of inhibiting telomerase activity in human breast cancer cells. These reagents will have potential clinical applications to breast cancer therapy and prevention.